



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Hence there are six sets of solutions:

$$x=0, y=0; x=-\frac{p}{m}\omega^3, y=-\frac{p}{m}\omega^2.$$

(2) Solution of the simultaneous equations

$$mx^4+py=0, my^3+px^3=0.$$

In addition to the solutions $x=y=0$, there are exactly nine sets of solutions

$$x=\varepsilon p^{4/9}m^{(-4)/9}, y=-\varepsilon^4 p^{7/9}m^{(-7)/9},$$

where ε is an arbitrary ninth root of unity.

165. Proposed by J. K. ELLWOOD, A. M., Principal of Colfax School, Pittsburg, Pa.

Solve $x^4-x=14$, by quadratics.

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

$$x^4-x=14 \text{ or } x^4-16=x-2.$$

$$\therefore x^2-4=\frac{x-2}{x^2+4}=\frac{x}{x^2+4}-\frac{2}{x^2+4}.$$

$$\therefore x^2-\frac{x}{x^2+4}+\frac{1}{4(x^2+4)^2}=4-\frac{2}{x^2+4}+\frac{1}{4(x^2+4)^2}.$$

$$\therefore x=2 \text{ or } x=-2+\frac{1}{x^2+4}.$$

$$\therefore x=2 \text{ or } x^3+2x^2+4x+7=0.$$

$$\therefore x=2 \text{ or } =-1\frac{1}{8} \text{ nearly, or } -\frac{1}{18}[1\pm\sqrt{(-843)}] \text{ nearly.}$$

166. Proposed by MARCUS BAKER, U. S. Geological Survey, Washington, D. C.

Solve

$$ax+by=2zx\dots(1).$$

$$cy+dz=2xy\dots(2).$$

$$ez+fx=2yz\dots(3).$$

Solution by the PROPOSER.

From (1), (2), and (3), respectively,

$$y=\frac{(2z-a)x}{b}=\frac{dz}{2x-c}=\frac{ez+fx}{2z},$$

whence

$$x(2x-c)(2z-a)=bdz\dots(4),$$

$$fx(2x-c)+ez(2x-c)=2dz^2\dots(5).$$

From (4), $z = \frac{ax(2x-c)}{2x(2x-c)-bd}$, which substituted in (5), gives, after reduction,

$$x^4 + \frac{1}{2f}(ae-2cf)x^3 + \frac{1}{4f}[fc^2 - a^2d - 2(ace + bdf)]x^2 \\ + \frac{1}{8f}[ac(ad+ce) + bd(2cf-ae)]x + \frac{1}{16f}bd(ace + bdf) = 0.$$

Similarly,

$$y^4 + \frac{1}{2b}(ac-2be)y^3 + \frac{1}{4b}[fc^2 - e^2b - 2(ace + bdf)]y^2 \\ + \frac{1}{8b}[ce(ae+cf) + df(2be-ac)]y + \frac{1}{16b}df(ace + bdf) = 0.$$

$$z^4 + \frac{1}{2d}(ce-2ad)z^3 + \frac{1}{4d}[a^2d - e^2b - 2(ace + bdf)]z^2 \\ + \frac{1}{8d}[ac(ae+be) + bf(2ad-ce)]z + \frac{1}{16d}bf(ace + bdf) = 0.$$

Also solved by LON C. WALKER.

GEOMETRY.

REMARKS ON NO. 187, GEOMETRY, BY J. R. HITT, GOSS, MISS.

There seems to be an error in (4) of Professor Zerr's demonstration of No. 187, Geometry. The result given is not correct. For $t = \frac{1}{\sqrt{2}}b\sin C\cos C$, $t_1 = \frac{1}{\sqrt{2}}b\cos C$, $t_2 = \frac{1}{\sqrt{2}}b\sin C$, from which it is seen that in general t^2 cannot equal t_1t_2 . It is also easily seen that if $t_1 : t = t : t_2$, then must $DI = b$, whereas DI cannot be $> \frac{1}{2}b$.

CALCULUS.

154. Proposed by B. R. DOWNER, Hopkinsville, Ky.

At the equinox, when the sun is on the celestial equator, a man starts driving on a perfectly level plain at six o'clock in the morning, and continues, going always from the sun, at the uniform rate of six miles per hour, until six o'clock in the evening. Required the path he will travel and the distance in a straight line from starting point to stopping point.